Lithologic description of cuttings (part A) from drill hole USBM-01, Ryan Gulch

Description	Depth interval feet (meters)
Quaternary	
Alluvium, not cemented, moderate-yellowish-brown, very fine grained (0.25 mm); grains mostly quartz; sparse biotite flakes	30-45 (9.1-13.7)
Alluvium, possible siltstone, moderate-yellowish-brown, very fine	
grained, same as above; contains 4 mm fragments of siltstone and	
sparse fragments of well-cemented sandstone (<2 mm) or quartzite(?) Tertiary	45-60 (13.7-18.3)
Uinta Formation	
Siltstone, pale-yellowish-brown to dark-yellowish-brown, very fine	
grained, very thinly laminated, well cemented with silica	60-75 (18.3-22.9)
Siltstone, predominately dark-yellowish-brown, very fine grained,	(10.3-22.9)
well-cemented; some laminae are grayish yellow; contains sparse	
very thinly laminated kerogenous marlstone	75-90 (22.9-27.4)
Siltstone, same as above, some dusky-yellowish-brown	90-105 (27.4-32.0)
Siltstone, predominately dusky-yellowish-brown, well-cemented	(=:::)
(some silica and calcite); laminae are dark yellowish brown	
and grayish yellow; contains sparse laminae of kerogenous	
marlstone; breaks with concoidal fracture	105-120 (32.0-36.6)
Siltstone, pale-yellowish-brown, massive ~90 percent, and dusky-	
Yeilowish-gray kerogenous marlstone; calcite cement Siltstone, same as above	120-135 (36.6-41.1)
distribution of the contract o	135-150 (41.1-45.7)
Cement; interval at base of cemented casing	150-165 (45.7-50.3)
Siltstone, dark-yellowish-brown; laminated with dark-gray silt-	(.3., 00.0)
stone; scattered (0.25 mm) flakes of biotite that equal 1 per-	
cent of rock; some parts have more dark-gray laminae; pyrite	
present in discrete large masses	165-180 (50.3-54:9)

Description	Depth interval feet (meters)
TertiaryContinued	•
Uinta FormationContinued	
Sandstone, dark-gray, fine- to medium-grained, and minor amounts	
of light-brown, calcite-cemented siltstone; no calcite in	
sandstone; 70 percent of the sample is sandstone; dusky-	
yellowish-brown, fine- to medium-grained sandstone; contains	
tiny reddish-brown hematite, minor amounts of biotite, and	
scattered fine grains of rounded frosted quartz	(54.9-59.4)
No sample	195-290 (59.4-88.4)
Siltstone, dusky-yellowish-brown, laminated with kerogenous	
marlstone; contains minor amount of quartzitic sandstone	
and pyrite fragments	290-305 (88.4-93.0)
Sandstone, olive-gray to dusky-yellowish-brown, fine- to medium-	
grained; contains fragments of siltstone and subrounded quartz,	
minor biotite flakes, some kerogenous marlstone	305-330 (93.0-100.6)
Sandstone, same as above but with more angular quartz fragments	330-345 (100.6-105.2)
Sandstone and some siltstone, light-olive-gray; sandstone is	
fine grained with minor calcite cement and contains 85-90	1 2 2
percent quartz, <5 percent pyrite; both are cemented (silica	
cement and minor calcite); from cuttings, unit appears to be	
very thin interbedding of siltstone and sandstone	345-360 (105.2-109.7)
Siltstone, dusky-yellowish-brown, well-cemented; contains	
scattered subrounded quartz and minor (<5 percent) pyrite	360-375 (109.7-114.3)
Sandstone, dusky-yellowish-brown, fine- to medium-grained sub-	•
angular quartz; contains pyrite; some siltstone appears to	
be poorly cemented	375-390 (114.3-118.9)
Siltstone, dark-yellowish-brown, very fine to fine-grained,	
well-cemented; contains minor sandstone (90 percent quartz);	
minor kerogenous marlstone stringers scattered throughout	390-405 (118.9-123.4)

Description	Depth interval feet (meters)
TertiaryContinued	F 3
Uinta FormationContinued	3.
Siltstone, dark-yellowish-brown to dusky yellowish-brown, mostly	
quartz; contains minor sandstone (90 percent quartz), abundant	P
pyrite, and minor kerogenous marlstone; all well cemented with	D # 258
silica and minor calcite	405-420 (123.4-128.0)
Siltstone and sandstone, pale-yellowish-brown to dusky-yellowish-	
brown, very fine to fine-grained, well-cemented; contains minor	
amounts of kerogenous marlstone; sandstone contains pyrite	420-435 (128.0-132.6)
Interbedded sandstone and siltstone, dark-yellowish-brown; 70	
percent sandstone, 30 percent siltstone; contains minor amounts	
of kerogenous marlstone and minor pyrite; unconsolidated	435-450 (132.6-137.2)
Interbedded siltstone, marlstone, and sandstone, dark-yellowish-	
brown to dusky-yellowish-brown; 40 percent siltstone, 40 per-	
cent marlstone, 20 percent coarse-grained sandstone; some	D)
kerogenous stringers; unconsolidated	450-465 (137.2-141.7)
Interbedded marlstone and siltstone, dark-yellowish-brown; 50	
percent marlstone, 50 percent unconsolidated siltstone	465-495 (141.7-150.9)
Same as above; possibly >60 percent marlstone	495-540 (150.9-164.6)
Laminated siltstone and kerogenous marlstone; siltstone (~ 50	
percent of sample) is pale yellowish brown with minor amounts	
of dark-yellowish-brown calcareous cement; kerogenous marl-	
stone (∿50 percent of sample) is dusky yellowish brown;	
contains minor amounts of frosted quartz	540-555 (164.6-169.2)
Same as above, with siltstone increasing to ${\sim}90$ percent	555-570 (169.2-173.7)
Same as above, with siltstone increasing to >95 percent	570-585 (173.7-178.3)
Laminated siltstone and kerogenous marlstone; siltstone (80 per-	a 6
cent) is mostly pale yellowish brown, some dark yellowish brown,	
contains calcareous cement; marlstone (20 percent) is medium	
gray	585-600 (178.3-182.9)

pth interval feet (meters)

5-420 4-128.0)

0-435 0-132.6)

i-450 i-137.2)

-465 -141.7)

495 150.9) 540 164.6)

555 69.2) 70 73.7) 85 78.3)

10 12.9)

Description	Depth interval feet (meters)
Tertiary Continued	
Uinta FormationContinued	* 1
Laminated siltstone and kerogenous marlstone: siltstone (90 per-	
cent) is dark yellowish brown, contains calcareous cement;	
dusky-yellowish-brown marlstone	
laminated kanaganawa mawlatawa a la say	(182.9-187.5)
Laminated kerogenous marlstone and siltstone; marlstone (90	
percent) is grayish black to olive black; siltstone (∿10 per-	
cent) is pale yellowish brown and contains calcareous cement	615-630 (187.5-192.0)
Laminated kerogenous marlstone and fine-grained sandstone with	
kerogenous streaks; marlstone (95 percent) is grayish black	
to olive black; sandstone (5 percent) is yellowish gray	630-645 (192.0-196.6)
Laminated kerogenous marlstone and sandstone; marlstone, 90	
percent; sandstone, 10 percent	645-660 (196.6-201.2)
Laminated kerogenous marlstone and sandstone; marlstone (90	(13010 20112)
Percent) is dusky yellowish brown and pale yellowish brown;	
sandstone (10 percent) is fine grained	660-675
	(201.2 - 205.7)
Kerogenous marlstone, dusky-yellowish-brown	675-690 (205.7-210.3)
Kerogenous marlstone, dusky-yellowish-brown; calcite fills	
fractures; trace of pale-yellowish-brown, fine-grained	
sandstone	690-700 (210.3-213.4)
Green River Formation	
Marlstone, kerogenous, dusky-yellowish-brown; trace of pale-	
yellowish-brown, fine-grained sandstone with kerogenous streaks	700-720 (213.4-219.5)
Marlstone, kerogenous as above, with calcite coatings on fracture	
surfaces	720-735 (219.5-224.0)
Marlstone, kerogenous as above, 90 percent; 10 percent grayish-	
orange-pink, fine- to medium-grained sandstone with calcite	-
cement; contains streaks of kerogen and sparse flakes of	
biotite	735-750 (224.0-228.6)
Bottom of rock-bit zone	

Description	Depth interval feet (meters)
Tertiary	
Green River Formation	5 Tu Har
	Committee of the commit
	•
on open fracture surfaces and in healed fractures.	
core ranges from competent to highly fractured	750-820.2
No sample	820.2-821.0 (250.0-250.2)
Oil shale, dusky-brown to dusky-yellowish-brown; calcite and nahcolite crystals on open and healed fractions.	Switch the of men.
	821.0-830.0
0.3	830.0-831.0 (253.0-253.3)
Oil shale, like unit at 821.0-830.0 feet (250.2-253.0 m);	
dusky yellowish brown is dominant color below the	
Marlstone, dusky-, dark-, and pale-yellowish-brown work	831.0-853.0 (253.3-260.0)
thinly laminated; calcite in open and healed fractures.	776 Jan
pyrite crystals on fractures in lowest 1 foot (30 cm); occasional vugs, some contain nahcolite	
Siltstone, interlaminated and bedded with many	(260.0-262.1)
Groove A	the second of the second of the second
Oil shale, pale-yellowish-brown and dark- and ducky walk	(262.1-264.6)
brown, very thinly laminated; contains pyrite crystals	of a mand by
No sample	- 868.0-869.0 (264.6-264.9)
Oil shale, like unit at 853.0-860.0 feet (260 0-262 1)	- 869.0-870.0 (264.9-265.2)
brecciated layer of oil shale in a silty matrix, with	* . 1
pyrite crystals at 875.0-875.6 feet (266.7-266.9 m)	- 870.0-886.3 (265.2-270.1)

interval eet ters)

Andrews of state of the state o

.2 .0) .0 .2)

0

)

Description	Depth interval feet (meters)
TertiaryContinued	
Green River FormationContinued	
Parachute Creek MemberContinued	
Siltstone, dark-yellowish-brown; contains streaks of kerogen	- 886.3-886.5 (270.1-270.2)
Oil shale, like unit at 853.0-860.0 feet (260.0-262.1 m)	- 886.5-889.3 (270.2-271.1)
Oil shale, dusky-yellowish-brown, laminated	- 889.3-891.4 (271.1-271.7)
Oil shale, like unit at 853.0-860.0 feet (260.0-262.1 m); pyrite	
crystals in lower 1 foot (30 cm)	(271.7-272.2)
No sample	893.0-895.0 (272.2-272.8)
Oil shale, dusky-yellowish-brown, laminated with dark-yellowish-	
brown oil shale; pyrite and calcite crystals disseminated	
throughout core; some discing of core	(272.8-275.5)
Siltstone, dusky-yellowish-brown	(275.5-275.7)
Oil shale, like unit at 895.0-904.0 feet (272.8-275.5 m)	(275.7-278.4)
No sample	913.5-915.0 (278.4-278.9)
Oil shale, dusky-yellowish-brown, thinly laminated with pale-	
yellowish-brown oil shale; calcite on healed fractures	915.0-928.2 (278.9-282.9)
Siltstone, pale-yellowish-brown; contains thin wavy beds of	
kerogenous material	928.2-928.4 (282.9-283.0)
Oil shale, like unit at 915.0-928.2 feet (278.9-282.9 m);	*
core shattered and disced; vuggy	928.4-929.0 (283.0-283.2)
Oil shale, dark-yellowish-brown, laminated with dusky-yellowish-	
brown oil shale; pale yellowish brown in lowest 1 foot (30 cm);	4.4
some laminae are wavy, most are flat and horizontal; pyrite	r es gil
and calcite crystals on open and healed fractures; minor	
feathering of some laminae	929.0-949.0 (283.2-289.3)
Gilsonite(?), laminated with oil shale and a white micro-	
crystalline mineral (noncalcareous)	- 949.0-949.8 (289.3-289.5)

randri orazola Mari Morrismo)	Description		Depth interval feet (meters)
TertiaryContinued			e Çerê di ave se
Green River Formation	Continued	28 165	rellinger a management
Parachute Creek Membe	rContinued	,	e et trippi de a servi
		283.2-289.3 m)	(289.5-290.1)
		sand-size matrix	(290.1-290.1+)
* **		283.2-289.3 m)	(290.1+-298.7)
		with dusky-yellowish-	
	precciated zones at 982		g r B t _e y a
	0.5 foot (15 cm) thick;		
	opear at 989.7 feet (301		
	eet (303.0 and 303.6 m);		de la companya de la
	s at 987.0 feet (300.8 m		
			(298.7-303.9)
A		aminated oil shale	997.0-997.8 (303.9-304.1)
	t at 980.0-997.0 feet (2		
	etween 1,000.0 and 1,006	0.0 feet (204.8	8 800m U. 20 a 90mm g
	cciated zones at 1,001.0		
	nd 1,011.0-1,014.0 feet		
		below 1,007.5 feet	
			(304.1-308.8)
Sandstone, medium-gr	ained		1,013.0-1,013.2±
Oil shale, like unit		98.7-303.9 m)	(308.8-308.8±)
on one of the unit	2 40 300.0 337.0 1000 (2		(308.8±-309.4)
Oil shale, grayish-l	olack to black		1,015.0-1,015.5 (309.4-309.5)
Oil shale, dark-yell	lowish-brown; interlamin	ated with dusky-	10 a.60
		and thin beds of	
siltstone mainly b	petween 1,016.0 and 1,01	9.1 feet (309.7-	a terrisi gela
310.6 m) and 1,022	2.0 and 1,023.0 feet (31	1.5-311.8 m);	and the second
fairly competent o	core		1,015.5-1,023.0 (309.5-311.8)
	to and the state of		
Δu-ij '			

nterval et ers)

7

9 1+)

0 7)

.2± ±)

0

0

Description	Depth interval feet (meters)
TertiaryContinued	
Green River FormationContinued	
Parachute Creek MemberContinued	х н
Oil shale, dark-yellowish-brown; interlaminated with minor	
amounts of very dusky red oil shale; contains several	
brecciated zones of oil-shale fragments in a sandy-silty	1,023.0-1,039.0
matrix	(311.8-316.7)
No sample	1,039.0-1,039.2 (316.7-316.7+)
Oil shale and clay(?); upper 5 feet (1.5 m) is dark-	
yellowish-brown oil shale containing vugs partially	
filled with nahcolite; lowest 4 feet (1.2 m) is pale-	
yellowish-brown clay containing a few 2-inch (5-cm)	
fragments of oil shale	1,039.2-1,048.2
	(316.7+-319.5) 1,048.2-1,052.6
No sample	(319.5-320.8)
Oil shale, light- to dusky-yellowish-brown; interlaminated	
with light-yellowish-brown siltstone	1,052.6-1,063.0 (320.8-324.0)
Oil shale, pale- to dusky-yellowish-brown, with minor beds	*
of pale-yellowish-brown tuff(?); leaching evident at 1,072	
and 1,083 feet (326.7 and 330.1 m); nahcolite in vugs at	
1,083 feet (330.1 m); core discs easily	1,063.0-1,083.5 (324.0-330.3)
No sample	1,083.5-1,087.0 (330.3-331.3)
Like unit at 1,063.0-1,083.5 feet (324.0-330.3 m); tuff(?)	
layers occur between 1,088.6 and 1,089.6 feet (331.8 and	
332.1 m)	1,087.0-1,093.0 (331.3-333.1)
Oil shale, pale- to dusky-yellowish-brown, thinly to very	
thinly laminated; calcite on fracture surfaces; sparse	
vugs as much as 2 inches (5 cm) in diameter contain minor	1 002 0 1 100 0
amounts of nahcolite	1,093.0-1,109.0 (333.1-338.0)
Oil shale, as above, interbedded with pale- to moderate-	
yellowish-brown siltstone beds at 1,109-1,110, 1,112, 1,117,	
1,120-1,121, and 1,125 feet (338.0-338.3, 338.9, 340.5, 341.4-	1 100 0 1 105 0
341.7, and 342.9 m)	1,109.0-1,125.0 (338.0-342.9)
25	Section 201

	15 1 e	Description	Depth interval feet (meters)
Tert	iaryContinued		
Gr	een River Formation	Continued	1. "0.7"
	Parachute Creek Men	berContinued	# # # # # # # # # # # # # # # # # # #
	Oil shale, like u	nit at 1,093.0-1,109.0 feet (333.1-338.0 m),	with
		f(?) bed at 1,149 feet (350.2 m); collapse	
	breccia zone at	1,142.0-1,142.6 feet (348.1-348.3 m) may ma	tch
		,176.5 feet (357.8-358.6 m) in drill hole	
	USBM-02A; core-	loss interval below may match collapse breco	ia
		eet (361.8 m) in USBM-02A	
			(342.9-351.2)
			1,152.2-1,155.2 (351.2-352.1)
		ellowish-brown, interlaminated with minor	
		ae of dark- and dusky-yellowish-brown to	
		shale; contains a few scattered vugs as wid cm), some containing minor amounts of	e
		some containing minor amounts of	1 155 0 1 167 0
			(352.1-355.7)
		ve, interlaminated with thin-bedded,	
		ish-brown siltstone; possible thin tuff	
		et (355.7 m)	1,167.0-1,172.5 (355.7-357.4)
		nit at 1,155.2-1,167.0 feet (352.1-355.7 m);	
		as at 1,172.5-1,174.5 feet (355.7-358.0 m)	
		78.0 feet (358.3-359.1 m); a thin, medium-	
		tiferous siltstone at 1,174.5 feet (358.0 m)	· · · · · · · · · · · · · · · · · · ·
		-1,174.5 feet (357.4-358.0 m) contains a	
		crystalline mineral; unit may correspond to	
		val at 1,201.8-1,209.5 feet (366.3-368.7 m) SBM-02A	
	in drift hore of	3011-0271	1,172.5-1,178.0 (357.4-359.1)
	Oil shale, dark-ye	ellowish-brown, very thinly laminated; thin,	(00,11,003,17)
		ish-brown siltstone layer at 1,181 feet	F. D. DERF W. F. F. S.
		rately well cemented; dusky-yellowish-brown	in Since Object of April
	to dusky-brown	oil shale at 1,185-1,189 feet (361.2-362.4 m);
	open vugs, as m	uch as 2 inches (5 cm) contain minor amounts	en Fa
		vidence of leaching at 1,197.4-1,198.0 feet	
		minor amounts of calcite and pyrite on	*44 .4 6
	fracture surface	es; unit is brittle and incompetent	1,178.0-1,203.9
			(359.1-366.9)

Lithologic description of core (part B) from drill hole USBM-01, $Ryan\ \textit{Gulch}\text{--}\texttt{Continued}$

** *** **** **** **** ****	
Description	Depth interval feet (meters)
TertiaryContinued	
Green River FormationContinued	
Parachute Creek MemberContinued	
No sample	1,203.9-1,207.2 (366.9-368.0)
Siltstone, pale-yellowish-brown and yellowish-gray, very thinly	2. 2
laminated; some is thin bedded; contains minor amounts of	
dark- to dusky-yellowish-brown, very finely laminated oil	
shale and marlstone(?); faults at 1,218 and 1,230 feet	
(371.2 and 374.9 m)the lower one has abundant calcite	
dispersed throughout zone	1,207.2-1,230.0 (368.0-374.9)
Marlstone, pale-yellowish-brown, yellowish-gray, and dark-	
Yellowish-brown, very thinly laminated to thin-bedded;	
contains minor amounts of dusky-yellowish-brown oil-shale	
laminae and thin siltstone stringers; vugs, 1/4 inch (1	
cm) in diameter, filled with calcite and minor amounts of	
nahcolite; calcite and pyrite on fracture surfaces; base	
of R-6 zone at 1,240 feet (378.0 m)	1,230.0-1,287.7 (374.9-392.5)
No sample	1,287.7-1,292.7 (392.5-394.0)
Siltstone, pale-yellowish-brown, yellowish-gray, and medium-	¥
gray, thinly bedded; contains minor amounts of marlstone	9.0
and oil shale; vugs (as wide as 3 in.) contain calcite and	121
nahcolite; pyrite and calcite on fracture surfaces;	t
collapse breccia or fault zone at 1,318-1,320.8 feet	
(401.7-402.6 m)	1,292.7-1,320.8 (394.0-402.6)
No sample; base of L-5 zone at 1,327 feet (403.6 m)	1,320.8-1,330.3 (402.6-405.5)
Oil shale, dusky-yellowish-brown and dusky-brown, very	
thinly laminated; contains minor amounts of siltstone	
and marlstone; evidence of leaching at 1,330.3-1,331.3	
feet (405.5-405.8 m); small blebs of nahcolite at	10 8 8 8 6
1,330.3-1,332.7 feet (405.5-406.2 m)	1,330.3-1,335.1 (405.5-406.9)
No sample	1,335.1-1,335.9 (406.9-407.2)

-1,203.9 -366.9)

pth interval feet (meters)

i.0-1,152.2 .9-351.2)

.2-1,155.2 .2-352.1)

.2-1,167.0 .1-355.7)

0-1,172.5 7-357.4)

ner off

5-1,178.0 1-359.1)

er ev	Description	Depth interval feet (meters)
TertiaryContinued		20 T
Green River Formation	onContinued	
Parachute Creek M		a asa ^{ari} a
	-yellowish-brown and yellowish-gray, very	and the second
	ted, platy; interlaminated with nahcolite	
		1,335.9-1,353.4 (407.2-412.5)
No sample		1,353.4-1,363.8 (412.5-415.7)
Oil shale, naho	olite, and halite, like unit at 1,335.9-	
1,353.4 feet	(407.2-412.5 m)	1,363.8-1,370.5 (415.7-417.7)
Oil shale, brow	nish-black and dusky-yellowish-brown, very	
thinly lamina	ted; contains rosettes of nahcolite	1,370.5-1,371.4 (417.7-418.0)
Oil shale, pale	-yellowish-brown and yellowish-gray, very	
thinly lamina	ted; interlaminated with pale-brown nahcol	ite
and minor amo	unts of halite; crystalline zone at 1,373.	1-
1,379.1 feet	(418.5-420.3 m); pitting along laminae in	this
crystalline z	one; calcite on some fracture surfaces	1,371.4-1,385.5 (418.0-422.3)
Oil shale, dusk	y-yellow-brown and brownish-black, very	
	ted; light-brown transparent nahcolite	
rosettes scat	tered throughout rock	1,385.5-1,393.8 (422.3-424.8)
	rate-brown, dark-yellowish-brown, and dusk	
	chinly laminated; minor laminae of marlstor	
	nt-brown nahcolite rosettes as much as 6 in	
(15.2 cm) in	diameter and minor halite beds	1,393.8-1,421.2 (424.8-433.2)
	erate-brown, moderate-yellowish-brown and c	
	own, very thinly laminated; intervals betwe	
SE SECRETARISE MASSES NO.	3.0, 1,451.0-1,477.0, 1,484.0-1,486.0, and	
50 (50)	9.0 feet (435.9-441.4, 442.3-450.2, 452.8-	
	53.3-453.8 m) contain nahcolite rosettes as	
Ø 38	ches (15.2 cm) in diameter	(433.2-453.8)
Halite, white,	translucent	1,489.0-1,492.0 (453.8-454.8)

Depth interval feet (meters)	Description	Depth interval feet (meters)
The second of the second secon	TertiaryContinued	
	Green River FormationContinued	* = 29
III waa	Parachute Creek MemberContinued	•
* 1.*	Oil shale, dark- and dusky-yellowish-brown, very thinly	.61
F10 1 2	laminated; with minor amounts of yellowish-gray and	
5.9-1,353.4	dusky-brown oil shale; moderate-brown translucent	
7.2-412.5)	crystalline nahcolite in beds, rosettes, and pods	
3.4-1,363.8 2.5-415.7)	scattered through unit	1,492.0-1.510.8
2.0 413.7)		(454.8-460.5)
3.8-1,370.5	Nahcolite, moderate-brown, translucent, and pale-yellow-	
5.7-417.7)	ish-brown, opaque	1,510.8-1,515.0
	Oil shale, like unit at 1,492.0-1,510.8 feet (454.8-	(460.5-461.8)
0.5-1,371.4	460.5 m); contains small amounts of olive-black oil	
7.7-418.0)	shale	1 515 0 1 504 0
		(461.8-464.5)
	Nahcolite, like unit at 1,510.8-1,515.0 feet (460.5-	
	461.8 m)	1,524.0-1,534.0
-4-1,385.5		(464.5-467.6)
.0-422.3)	Oil shale, dusky- to dark-yellowish-brown, very thinly	
k k	laminated; contains pods and irregular lenses of	
Section 2	moderate-brown, translucent, crystalline nahcolite	1,534.0-1,549.0 (467.6-472.1)
.5-1,393.8	Nahcolite, pale-yellowish-brown, opaque, crystalline	
.3-424.8)		(472.1-473.7)
The same was	Oil shale, dusky-yellowish-brown, dusky-brown, brownish-	
413	black, and minor dark-yellowish-brown, very thinly	
	laminated; contains a 0.8-foot-thick (24-cm) nahcolite	
.8-1,421.2 .8-433.2)	bed at 1,575.0 feet (480.1 m)	1,554.0-1,587.0
100.2)	Nahcolite white to pale vellowich have a hadder	(473.7-483.7)
and the second	Nahcolite, white to pale-yellowish-brown, bedded, opaque,	
and the state of t	crystalline; interbedded with laminae of oil shale	1,587.0-1,598.3 (483.7-487.2)
and the state of t	Oil shale, like unit at 1,554-1,587 feet (473.7-483.7 m)	
2.10		
2-1,489.0	No sample	1,599.3-1,600.8
2-453.8)	011 Shale, like unit at 1 554-1 507 foot (472 7 402 7 -)	(487.5-487.9)
0-1,492.0	Oil shale, like unit at 1,554-1,587 feet (473.7-483.7 m)	1,600.8-1,603.8 (487.9-488.8)
8-454.8)		

Description	Depth interval feet (meters)
TertiaryContinued	4 6 1
Green River FormationContinued	
Parachute Creek MemberContinued	7.3
Halite, bedded	1,603.8-1,604.3 (488.8-489.0)
Nahcolite and oil shale, interbedded	1,604.3-1,610.0 (489.0-490.1)
Oil shale, like unit at 1,554-1,587 feet (473.7-483.7 m)	1,610.0-1,616.0 (490.1-492.6)
Nahcolite and oil shale, interbedded	1,616.0-1,620.5 (492.6-493.9)
Oil shale, like unit at 1,554-1,587 feet (473.7-483.7 m)	1,620.5-1,622.6 (493.9-494.6)
Nahcolite and oil shale, interbedded	1,622.6-1,629.0 (494.6-496.5)
Halite, bedded	1,629.0-1,629.5 (496.5-496.7)
Nahcolite and oil shale, interbedded	1,629.5-1,632.0 (496.7-497.4)
No sample (loss due to solubility of salts)	1,632.0-1,639.0 (497.4-499.6)
Oil shale, dusky-yellowish-brown, dark-yellowish-brown and	
minor light-brown, very thinly laminated; interbedded with	2
halite zones, as much as 0.5 foot (15 cm) thick, and	
moderate-brown, crystalline, translucent nahcolite beds	
and rosettes; some nahcolite is pale yellowish brown	1,639.0-1,660.0 (499.6-506.0)
Oil shale, as above	1,660.0-1,667.7 (506.0-508.3)
No sample	1,667.7-1,669.0 (508.3-508.7)
Oil shale, like unit at 1,639.0-1,660.0 feet (499.6-506.0 m); contains scattered nahcolite rosettes; base of R-5 zone at	4.4
1,680.9 feet (512.3 m)	1,669.0-1,680.9 (508.7-512.3)
Nahcolite, dusky- to dark-yellowish-brown; contains a halite	
stringer at 1,682 feet (512.7 m)	1,680.9-1,686.9 (512.3-514.2)
No sample	1,686.9-1,699.0 (514.2-517.9)
Nahcolite, like unit at 1,680.9-1,686.9 feet (512.3-514.2 m)	
N	

epth interval feet (meters)

8-1,604.3 8-489.0)

3-1,610.0 0-490.1)

0-1,616.0 1-492.6)

)-1,620.5 i-493.9)

-1,622.6 -494.6)

-1,629.0 -496.5) -1,629.5 496.7) 1,632.0 497.4)

,639.0 199.6)

660.0 667.7 8.3) 669.0 3.7)

80.9 .3)

9.0 9) 2.5 9)

Description	Depth interval feet (meters)
TertiaryContinued	
Green River FromationContinued	
Parachute Creek MemberContinued	
Oil shale, dusky-yellowish-brown, with minor amounts of pale-	
yellowish-brown, very thinly laminated oil shale; moderate-	
brown, translucent crystalline nahcolite rosettes as much	
as 3 inches (8 cm) in diameter scattered throughout core	1,702.5-1,715.0 (518.9-522.7)
Nahcolite, pale-yellowish-brown, oapque, crystalline,	2 E 2
thin-bedded	1,715.0-1,717.0 (522.7-523.3)
Oil shale, like unit at 1,702.5-1,715.0 feet (518.9-522.7 m);	
contains some brownish-black oil shale	(523.3-526.7)
No sample	(526.7 - 527.0)
Oil shale, like unit at 1,717.0-1,727.9 feet (523.3-526.7)	1,729.0-1,734.5 (527.0-528.7)
Nahcolite, moderate-brown, translucent, crystalline, thin-	
bedded; with minor amounts of oil shale	1,734.5-1,737.0 (528.7-529.4)
Oil shale, like unit at 1,717.0-1,727.9 feet (523.3-526.7 m)	1,737.0-1,739.0 (529.4-530.0)
Nahcolite, like unit at 1,734.5-1,737.0 feet (528.7-529.4 m);	
base of L-4 zone at 1,742 feet (531.0 m)	1,739.0-1,742.0 (530.0-531.0)
Oil shale, like unit at 1,717.0-1,727.9 feet (523.3-526.7 m);	
contains a 4-inch-thick (10-cm) bed of nahcolite and halite	
at 1,750.9 feet (533.7 m)	1,742.0-1,759.0 (531.0-536.1)
Nahcolite, pale-yellowish-brown and moderate-brown, opaque to	
translucent, thin-bedded; with minor oil shale	1,759.0-1,768.5 (536.1-539.0)
Oil shale, dark-yellowish-brown and dusky-yellowish-brown,	
very thinly laminated	1,768.5-1,777.5 (539.0-541.8)
Nahcolite, like unit at 1,759.0-1,768.5 feet (536.1-539.0 m)	
Oil shale, like unit at 1,768.5-1,777.5 feet (539.0-541.8 m)	1,779.5-1,783.0 (542.4-543.5)

Description	Depth interva feet (meters)
TertiaryContinued	(110 001 3)
Green River FormationContinued	
Parachute Creek MemberContinued	
Nahcolite, like unit at 1,759.0-1,768.5 feet (536.1-539.0 m)	
No sample	1,783.0-1,788.2 (543.5-545.0)
Oil shale, like unit at 1,768.5-1,777.5 feet (539.0-541.8 m);	7,788.2-1,789.0 (545.0-545.3)
Tosettes, as wide as A inches (10	
lite beds scattered throughout interval	111
No sample	1,789.0-1,818.9 (545.3-554.4)
017	1,818.9-1,819.4
Oil shale, dusky-yellowish-brown, contains rosettes of nahco-	(554.4-554.6)
lite as much as 9 inches (23 cm) in diameter	- 1 910 4 1 000 4
Nahcolite, bedded and rosettes	1,819.4-1,828.4 (554.6-557.3)
	.,020.7-1,030.0
Oil shale, nahcolite, and minor marlstone, interbedded and	(557.3-557.8)
rammated, pare-yellowish-brown to dusky-vellowish by	
rosettes as much as 10 inches (25 cm) in diameters balds	S 2 197
restriction tractures at 1.834 and 1.946 feet 1550	
and 562.7 m) Oil shale, dusky-vellowich-brown land	- 1,830.0-1,850.0 (557.8-563.9)
Oil shale, dusky-yellowish-brown, laminated with dusky- to dark-	1.5 %
yellowish-brown marlstone; nahcolite rosettes as much as 6 inches (15 cm) in diameter every foot or two; fine- to	
medium-grained nahoolite had at 1 070 a core	
medium-grained nahcolite bed at 1,873-1,873.5 feet (570.9-	
571.0 m), and a 0.1-foot-thick (3-cm) bed at 1,853.8 feet (565.0 m)	
No sample	
Nahcolite, interbedded with dark-yellowish-brown and minor	(571.2-572.7)
addry-yellowish-brown maristone(?): nabcolite	
	Commence of the Commence of th
stone rayer at 1,887.5-1,887.7 feet (575 3-575 4-1)	
solite losettes scattered throughout lower bals as	
(1,893-1,907.4 feet (577.0-581.4 m)\	
(1,893-1,907.4 feet (577.0-581.4 m))	1,879.0-1,907.4 (572.7-581.4)

terval t rs)

8.2 0) 9.0 3)

8.9 4)

9.4 5)

3.4

1.0

.0

Description	Depth interval feet (meters)
TertiaryContinued	9
Green River FormationContinued	
Parachute Creek MemberContinued	6
Oil shale, dark-yellowish-brown, thinly laminated; minor laminae	
are dusky yellowish brown; bedded opaque nahcolite at 1,908.2-	
1,908.4 and 1,921.3-1,921.5 feet (581.6-581.7 and 585.6-585.7	
m); nahcolite rosettes as much as 7 inches (18 cm) in diameter,	
most are 3-5 inches (8-15 cm); eroded rosettes cause core to	
break in 0.5-to 2-foot (15- to 60-cm) lengths	1,907.4-1,937.2 (581.4-590.5)
No sample	1,937.2-1,937.4 (590.5-590.5+)
Oil shale, dark- to dusky-yellowish-brown, thinly laminated;	
thin interbeds of nahcolite at 1,943, 1,960, 1,961.5, 1,969.8-	
1,970.2, and 1,972.3-1,972.9 feet (592.2, 597.4, 597.8, 600.4-	
600.5, and 601.2-601.3 m); nahcolite rosettes between 1,946	
and 1,967 feet (593.1 and 599.5 m) and scattered between	5
1,968 and 1,997 feet (599.8 and 608.7 m); abundant dolomite	*
lenses as long as 1 inch (3 cm) and as high as 1/8 inch	
(0.3 cm) throughout oil shale; "60 bed" at 1,988-1,992 feet	
(605.9-607.2 m); base of R-4 zone at 1,997 feet (608.7 m)	1,937.4-1,997.0 (590.5+-608.7)
Marlstone(?), dark-yellowish-brown, laminated with minor	
amounts of dusky-yellowish-brown oil shale; unit contains	
thin-bedded nahcolite at 2,002.0,2,004.6, 2,017.6, and	
2,020.6 feet (610.2, 611.0, 615.0, and 615.9 m); nahcolite	
rosettes 1-3 inches (3-8 cm) in diameter below 2,015 feet	
(614.2 m); bottom of L-3 zone at 2,026 feet (617.5 m)	1,997.0-2,025.9 (608.7-617.5)
No sample	2,025.9-2,027.1 (617.5-617.9)
Oil shale, dark-yellowish-brown, interlaminated and inter-	
bedded with bedded nahcolite, ranging from laminae to 3	
inches (8 cm) thick; unit contains pale-yellowish-brown	
lenses of dolomite as much as 2 inches (5 cm) long (most	. 163
are 1/4-1 in. (1-3 cm)); rock is 50 percent or more nah-	
colite; rosettes of nahcolite as much as 1.5 feet (45 cm)	
in diameter	2,027.1-2,085.7 (617.9-635.7)

Description	Depth interval feet (meters)
TertiaryContinued	
Green River FormationContinued	est to end of the
No sample	- 2,085.7-2,087.0 (635.7-636.1)
Oil shale, like unit at 2,027.1-2,085.7 feet (617.9-635.7 m),	
with nahcolite beds as much as 6 inches (15 cm) thick	- 2,087.0-2,094.0 (636.1-638.3)
Oil shale, dark- to dusky-yellowish-brown, containing	
nahcolite rosettes	(638.3-645.0)
No sample	(645.3-647.1)
Nahcolite, dark-yellowish-brown, bedded	(645.3-647.1)
Oil shale, dusky- to dark-yellowish-brown; unit contains opaque	A Section of the Sect
and translucent brown nahcolite rosettes as much as 1 foot	S 45 000
	and the same
yellow-brown lenses of dolomite below 2,137 feet (651.4 m)	/ 647 7 674 61
No sample	- 2,146.5-2,147.5 (654.3-654.6)
The state of the s	- (10 mm - 10
dark-yellowish-brown marlstone; upper 3 feet (90 cm) is interbedded marlstone, nahcolite, and nahcolite rosettes;	e e di Tanagara si
Dase 01 K-3 2011e	(654.6-659.3)
Marlstone(?) dark-yellowish-brown, thinly laminated	(659.3-663.1)
No sample	- 2,175.5-2,177.5 (663.1-663.7)
Marlstone(?), like unit at 2,163-2,175.5 feet (659.3-663.1 m)	
ကြောင်းသည်။ သို့ ကြောင်းသည်။ သည်သည်။ သ	(663.7-664.6)
Oil shale, dusky- and dark-yellowish-brown, thinly laminated	(664.6-667.2)
Nancolite, bedded	2,189.0-2,195.0
The state of the s	or the property

th interval feet (meters)

-2,087.0 -636.1)

-2,094.0 -638.3)

-2,116.3 -645.0)

-2,117.0 -647.1) -2,123.0 -647.1)

-2,146.5 -654.3) -2,147.5 -654.6)

-2,163.0 -659.3) -2,175.5 -663.1) -2,177.5 -663.7)

-2,180.5 -664.6)

-2,189.0 -667.2)

-2,195.0 -669.0)

* * · · ·

Description	Depth interval feet (meters)
TertiaryContinued	
Green River FormationContinued	
Parachute Creek MemberContinued	8 2 2 E 45
Oil shale, dark- to dusky-yellowish-brown, laminated; contains	
minor amounts of marlstone(?) and bedded nahcolite; nahcolite	4,54
rosettes as large as 8 inches (20 cm) (most are 2-4 in. (5-8	k #24
cm) scattered throughout core; dolomite lenses in upper 32	. 2
feet (9.8 cm)	- 2,195.0-2,245.0 (669.0-684.3)
Marlstone(?), dark-yellowish-brown, laminated; contains minor	
amounts of nahcolite	(684.3-687.1)
No sample	-,201.0 2,200.0
Oil shale, dusky- to dark-yellowish-brown, laminated	(687.1-688.2) - 2,258.0-2,268.0 (688.2-691.3)
Oil shale(?), dark-yellowish-brown, laminated; contains	(000:2-031:3)
dolomite lenses 1/8-1/2 inch (0.3-1 cm) long	- 2,268.0-2,281.0 (691.3-695.2)
Garden Gulch Member	
No sample. Blue marker at 2,284 feet (696.2 m); base of R-2	
Oil shale dusty to deal and an analysis of the state of t	- 2,281.0-2,286.0 (695.2-696.8)
Oil shale, dusky- to dark-yellowish-brown, laminated; dolo-	
Oil shale dusky-vellow brown laminated with the dusky brown laminate	2,286.9-2,290.5 (696.8-698.1)
Oil shale, dusky-yellow-brown, laminated; unit has what appear	
to be slickensides on horizontal fracture surfaces	2,290.5-2,299.0 (698.1-700.7)
Interbedded shale, marlstone, and a white bedded material;	
shale is bluish gray from 2,305 to 2,310 feet (702.6-704.1 m); marlstone is dusky yellowish brown, laminated to thin	
bedded; white material between 2,299 and 2,305 feet (700.7	
and 702.6 m) is in layers less than 0.1 foot (3 cm) thick;	
core discs easily; base of L-1 zone	
20119, 5036 01 1-1 20118	2,299.0-2,310.0 (700.7-704.1)

and the second	Description	E 58 :	**************************************	Depth interval feet (meters)
TertiaryContinued	•			34. " 1 3V "
Green River Formatio	nContinued		-,	
Garden Gulch Membe				
Marlstone oil s	hale, and shale, inter	laminated and	inter-	
bedded dark-	to dusky-yellowish-bro	wn; whitish-gra	ay,	
very thin heds	between 2,344.5-2,34	6.0 feet (714.	6- :	
715 1 m): foss	sils in 2,326-2,327-foo	t (709.0-709.3	-m)	
intorval may h	pe fish vertabrae; core	discs easily,		F. 51.5 mm (51.5
shattered in n	many places			2,310.0-2,382.0
		and the same of th		1
No sample, drill	led with rock bit			2,382.0-2,382.5 (726.0-726.2)
I to take stary				
., 337, 54, 63, (4,,112, 1,528)				2,382.5 (726.2)
election (1:4.4 Visite access			াৰ্ট , ভব	
	24,74,72	, se and the		e de la financia de la companya de l
7.85. 7 1, 40.5 168			at details	
	de man and a f	\$ 127 1 1 117		- 1.
a Bala da Tina. Talebah Maj	till sammen i sammen			
	er Fig. Wirth		and process	e da en la sistema de la composición della compo
			7 a 03 303 61 0 039	
1	46 State 2014		si i tagi tarih k	and the state
), and the second			Programme Comment	
	Ar walne	3000 - C	in a staff to a	1. 1. 1. 50 to 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
	1 n - 3.5.,		2.00	
	a table and the	140	e ger e area g	d the same
		11 11 11 11 11 11		. 21
			المعارة المعاري	A
9.2 3.3 5 5 7 5			. 1 . 34 al al .1	

Table | __ Lithologic description of core from drill hole USBM-01

Description	Interval thickness ft (m)
Core:	
Oil shale, dusky-yellowish-brown, laminated with	3.
minor amounts of dark-gray and grayish-brown oil	. 1
shale. Calcite and nahcolite crystals on open	
fracture surfaces and in healed fractures. Core	
ranges from competent to highly fractured	750-820.2 (228.6-250.0)
No recovery	820.2-821.0 (250.0-250.2)
Oil shale, dusky-brown to dusky-yellowish-brown.	
Calcite and nahcolite crystals on open and	
healed fracture surfaces. At 825.7-825.9 ft	
(251.7-251.7+ m) is a breccia of clay and oil	
shale fragments	821.0-830.0 (250.2-253.0)
Oil shale, grayish-black	830.0-831.0 (253.0-253.3)
Oil shale, like unit at 821.0-830.0 ft (250.2-253.0 m).	
Below 838 ft the laminae feather out across core.	
Dusky-yellowish-brown is dominant color below	
844 ft (257.3 m)	831.0-853.0 (253.3-260.0)
Marlstone, dusky-, dark-, and pale-yellowish-brown,	
very thinly laminated. Calcite in open and	
healed fractures, pyrite crystals on fractures	
in lower 1 ft (30 cm). Occasional vugs with	
some containing nahcolite	853.0-860.0 (260.0-262.1)

Description	Interval thickness ft (m)
CoreContinued	
Siltstone, interlaminated and bedded with marlstone.	
A Groove	860.0-868.0 (262.1-264.6)
Oil shale, pale-yellowish-brown with dark- and	
dusky-yellowish-brown, very thinly laminated	
pyrite crystals throughout unit	868.0-869.0 (264.6-269.4)
No recovery	869.0-870.0 (264.4-265.2)
Oil shale, like unit at $853.0-860.0$ ft ($260.0-262.1$ m).	
Brecciated layer of oil shale in a silty matrix	
with pyrite crystals 875.0-875.6 ft	
(266.7-266.9 m)	870.0-886.3 (265.2-270.1)
Siltstone, dark-yellowish-brown, containing	,
kerogenous streaks	886.3-886.5 (270.1-270.2)
Oil shale, like unit at 853.0-860.0 ft	
(260.0-262.1 m)	886.5-889.3 (270.2-271.1)
Oil shale, dusky-yellowish-brown, laminated	889.3-891.4 (271.1-271.7)
Oil shale, like unit at 853.0-860.0 ft (260.0-262.1 m).	
Pyrite crystals in lower 1 ft (30 cm)	891.4-893.0 (271.7-272.2)
No recovery. May be mahogany marker	893.0-895.0 (272.2-272.8)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, dusky-yellowish-brown laminated with dark	
yellowish brown. Pyrite and calcite crystals	
disseminated throughout core. Some discing of	
core. May be mahogany bed	895.0-904.0 (272.8-275.5)
Siltstone, dusky-yellowish-brown	904.0-904.6 (275.5-275.7)
Oil shale, like unit at 895.0-904.0 ft	*
(272.8-275.5 m)	904.6-913.5 (275.7-278.4)
No recovery	913.5-915.0 (278.4-278.9)
Oil shale, dusky-yellowish-brown, thinly laminated with	
pale yellowish brown. Calcite on healed fractures-	915.0-928.2 (278.9-282.9)
Siltstone, pale-yellowish-brown, contains thin wavy	•
beds of kerogenous material	928.2-928.4 (282.9-283.0)
Oil shale, like unit at 915.0-928.2 ft (278.9-282.9 m),	
core shattered and disced, vuggy	928.4-929.0 (283.0-283.2)
Oil shale, dark-yellowish-brown, laminated with dusky	
yellowish brown, pale yellowish brown in lower 1 ft	
(30 cm). Some laminae are wavy, most are flat and	
horizontal. Pyrite and calcite crystals on open	
and healed fractures. Minor feathering of some	,
laminae	- 929.0-949.0 (283.2-289.3)

Description	Interval thickness ft (m)
CoreContinued	
Gilsonite(?), laminated with oil shale and a white	
microcrystalline mineral (noncalcareous)	949.0-949.8 (289.3-289.5)
Oil shale, like unit at 929.0-949.0 ft	
(283.2-289.3 m)	949.8-951.7 (289.5-290.1)
Brecciated zone, oil shale with silt- and sand-size	e e
matrix	951.7-951.9 (290.1-290.1+)
Oil shale, like unit at 929.0-949.0 ft	*
(283.2-289.3 m)	951.9-980.0 (290.1+-298.7)
Oil shale, dark-yellowish-brown laminated with dusky	
yellowish brown. Brecciated zones at 982 and 986	
ft (299.3 and 300.5 m) each .5 ft (15 cm) thick.	,
Open and partially open vugs first appear at	
989.7 ft (301.7 m) with others at 994.0 and	
996.0 ft (303.0 and 303.6 m). Dusky-yellowish-	
brown color predominates 987.0 ft (300.8 m).	
Pyrite and calcite crystals on fracture surfaces	980.0-997.0 (298.7-303.9)
Sandy-silty layer interbedded with thinly laminated	
oil shale	997.0-997.8 (303.9-304.1)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, like unit at 980.0-997.0 ft (298.7-303.9 m).	
Discing of core between 1,000.0 and 1,006.0 ft	
(204.8 and 306.6 m). Brecciated zones at	
1,001.0-1,001.3 ft (305.1-305.2 m) and	
1,011.0-1,014.0 ft (308.2-309.1 m). Dusky-	
yellowish-brown color predominates below 1,007.5 ft	-
(307.1 m)	997.8-1,013.0 (304.1-308.8)
Sandstone, medium grained	1,013.0-1,013.2 ⁺ (308.8-308.8+)
Oil shale, like unit at 980.0-997.0 ft (298.7-303.9 m)	- 1,013.2±-1,015.0 (308.8+-309.4)
Oil shale, grayish-black to black	- 1,015.0-1,015.5 (309.4-309.5)
Oil shale, dark-yellowish-brown interlaminated with	
dusky yellowish brown, lenses, pods, and thin beds	
of siltstone mainly between 1,016.0-1,019.1 ft	
(309.7-310.6 m) and $1,022.0-1,023.0 ft$	
(311.5-311.8 m). Fairly competent core	- 1,015.5-1,023.0 (309.5-311.8)
Oil shale, dark-yellowish-brown interlaminated with	
minor amounts of very dusky red. Contains several	
brecciated zones of oil shale fragments in a	
sandy-silty matrix	- 1,023.0-1,039.0 (311.8-316.7)

Description		thickness (m)
CoreContinued ,		
No recovery	1,039.0-1 (316.7-3	
Oil shale and clay(?). Upper 5 ft (1.5 m) is dark-		
yellowish-brown oil shale with vugs partially		
filled with nahcolite. Lower 4 ft (1.2 m) is		
pale-yellowish-brown clay containing a few		
2-inch (5 cm) fragments of oil shale	1,039.2-1 (316.7+-	•
No recovery	1,048.2-1 (319.5-3	
Oil shale, light- to dusky-yellowish-brown inter-		
laminated with light-yellowish-brown siltstone	1,052.6-1 (320.8-3	
Oil shale, pale- to dusky-yellowish-brown with minor		
thin beds of pale-yellowish-brown tuff(?).		
Leaching evident at 1,072 and 1,083 ft (326.7	•	
and 330.1 m), nahcolite in vugs at 1,083 ft		
(330.1 m). Core discs easily	1,063.0-1 (324.0-3	
No recovery	1,083.5-1 (330.3-3	. D
Like unit at 1,063.0-1,083.5 ft (324.0-330.3 m).		
Tuff(?) layers occur between 1,088.6 and		* ,
1,089.6 ft (331.8 and 332.1 m)	1,087.0-1 (331.3-3	

	2 2 2
Description	Interval thickness ft (m)
CoreContinued ,	
Oil shale, pale- to dusky-yellowish-brown, thinly to	
very thinly laminated. Calcite on fracture sur-	-
faces, sparse vugs up to 2 inches (5 cm) in	
diameter contain minor amounts of nahcolite	1,093.0-1,109.0 (333.1-338.0)
Oil shale, as above, interbedded with pale- to	
moderate-yellowish-brown siltstone beds at	
1,109-1,110, 1,112, 1,117, 1,120-1,121, and	
1,125 ft (338.0-338.3, 338.9, 340.5, 341.4-341.7,	
and 342.9 m)	1,109.0-1,125.0 (338.0-342.9)
Oil shale, like unit at 1,093.0-1,109.0 ft	
(333.1-338.0 m), with a very thin $tuff(?)$ bed	
at 1,149 ft (350.2 m). Collapse breccia zone at	
1,142.0-1,142.6 ft (348.1-348.3 m) may match zone	
at 1,174-1,176.5 ft (357.8-358.6 m) in drill hole	
USBM-02A. Core loss interval below may match	• ,
collapse breccia zone at 1,187 ft (361.8 m) in	
02A	1,125.0-1,152.2 (342.9-351.2)
No recovery	1,152.2-1,155.2 (351.2-352.1)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, pale-yellowish-brown, interlaminated,	
with minor very thin laminae of dark- and dusky-	
yellowish-brown to olive-black oil shale. A	
few scattered vugs up to .75 inch (2 cm), some	
containing minor amounts of nahcolite	1,155.2-1,167.0 (352.1-355.7)
Oil shale, as above, interlaminated with thin bedded	
moderate yellowish-brown siltstone. Possible	
thin tuff bed at 1,167 ft (355.7 m)	1,167.0-1,172.5 (355.7-357.4)
Oil shale like unit at 1,155.2-1,167.0 ft	
(352.1-355.7 m). Collapse breccias at	
1,172.5-1,174.5 ft (355.7-358.0) and	
1,175.4-1,178.0 ft (358.3-359.1 m). A thin	ï
medium light gray pyritiferous siltstone at	×
1,174.5 ft (358.0 m). Zone at 1,172.5-1,174.5 ft	
(357.4-358.0 m) contains a layer of white crystal-	
line mineral. Unit may correspond to core loss	
interval 1,201.8-1,209.5 ft (366.3-368.7) in	*
drill hole USBM-02A	1,172.5-1,178.0 (357.4-359.1)

Description

Interval thickness ft (m)

Core--Continued

Oil shale, dark-yellowish-brown, very thinly laminated. Thin moderate-yellowish-brown siltstone layer at 1,181 ft (360.0 m), moderately well cemented. Dusky-yellowish-brown to dusky-brown oil shale at 1,185-1,189 ft (361.2-362.4 m). Open vugs, up to 2 inches (5 cm) in diameter. containing minor amounts of nahcolite. Evidence of leaching at 1,197.4-1,198.0 ft (365.0-365.2 m). Minor amounts of calcite and pyrite on fracture surfaces. Unit is brittle and incompetent-----1,178.0-1,203.9 (359.1 - 366.9)No recovery-----1,203.9-1,207.2 (366.9 - 368.0)Siltstone, pale-yellowish-brown and yellowish-gray, very thinly laminated, minor amounts are thin bedded; minor amounts of dark to dusky yellowish brown very finely laminated oil shale and marl-

> 1,207.2-1,230.0 (368.0-374.9)

stone(?). Faults at 1,218 and 1,230 ft (371.2

and 374.9 m); the lower one has abundant calcite

dispersed throughout zone-----

Description	Interval thickness ft (m)
CoreContinued	
Marlstone, pale-yellowish-brown, yellowish-gray, and	
dark-yellowish-brown; very thinly laminated to	
thin bedded. Minor amounts of dusky-yellowish-	
brown oil shale laminae. Thin siltstone	
stringers. Vugs, 1/4-inch (1 cm), filled with	
calcite and minor amounts of nahcolite, calcite	*.
and pyrite on fracture surfaces. Base of R-6	
zone at 1,240 ft (378.0 m)	1,230.0-1,287.7 (374.9-392.5)
No recovery	1,287.7-1,292.7 (392.5-394.0)
Siltstone, pale-yellowish-brown, yellowish-gray, and	
medium-gray, thinly bedded; minor amounts of	
marlstone and oil shale. Vugs (up to 3") contain	
calcite and nahcolite, pyrite and calcite on	•
fracture surfaces. Collapse breccia or fault	
zone at 1,318-1,320.8 ft (401.7-402.6 m)	1,292.7-1,320.8 (394.0-402.6)
No recovery. Base of L-5 zone at 1,324 ft (403.6 m)	1,320.8-1,330.3 (402.6-405.5)
Oil shale, dusky-yellowish-brown and dusky brown;	
very thinly laminated; minor amounts of siltstone	
and marlstone; evidence of leaching at 1,330.3-	
1,331.3 ft (405.5-405.8 m); small blebs of	
nahcolite 1,330.3-1,332.7 ft (405.5-406.2 m)	1,330.3-1,335.1 (405.5-406.9)

Description		thickness (m)
CoreContinued		T _a
No recovery	1,335.1-3	1,335.9 407.2) -
Oil shale, pale-yellowish-brown and yellowish-gray,		
very thinly laminated, platey; interlaminated		
with nahcolite and halite	1,335.9-3	
No recovery	1,353.4-3	
Oil shale, nahcolite, and halite like unit at		
1,335.9-1,353.4 ft (407.2-412.5 m)	1,363.8-3	100 · 100 ·
Oil shale, brownish-black and dusky-yellowish-brown;		
very thinly laminated; rosettes of nahcolite	1,370.5-2	
Oil shale, pale-yellowish-brown and yellowish-gray,		
very thinly laminated; interlaminated with pale	•	
brown nahcolite and minor amounts of halite,		
crystalline zone 1,373.1-1,379.1 ft (418.5-420.3 m).		
Pitting along laminae in this zone; calcite on		
some fracture surfaces	1,371.4-3	
Oil shale, dusky-yellow-brown and brownish-black,		
very thinly laminated, light brown transparent		
nahcolite rosettes scattered throughout rock	1,385.5-3	

	Description		thickness (m)
Core-	-Continued		
0i1	shale, moderate-brown, dark-yellowish-brown and		
	dusky-brown, very thinly laminated; minor laminae		
	of marlstone. Light-brown nahcolite rosettes to		
	6 inches in diameter and minor halite beds	1,393.8-	-1,421.2 -433.2)
0i1	shale, moderate-brown, moderate-yellowish-brown		-
	and dark-yellowish-brown, very thinly laminated.		
	Intervals between 1,416.5-1,421.5, 1,430-1,448,		
	1,451-1,477, 1,484-1,486 and 1,487.5-1,489 ft		
	(431.7-441.4, 435.9-441.4, 442.3-450.2,		
	452.8-452.9, and 453.4-453.8 m) contain		
	nahcolite rosettes up to 6 inches (15 cm) in		
	diameter	1,421.2-(433.2-	
Ha1	ite, white, translucent	1,489.0- (453.8-	
0i1	shale, dark- and dusky-yellowish-brown, with minor		
	amounts of yellowish gray and dusky brown, very		
	thinly laminated. Moderate brown translucent		
	crystalline nahcolite in beds, rosettes and pods		
	scattered through unit	1,492.0- (454.8-	
Nah	colite, moderate brown translucent and pale-		
	yellowish-brown opaque	1,510.8-	

Description		thickness (m)
CoreContinued		
Oil shale like unit at 1,492.0-1,510.8 ft		
(454.8-460.5 m) amounts of olive-black oil shale	1,515.0-1,524.0 (461.8-464.5)	
Nahcolite, like unit at 1,510.8-1,515.0 ft		
(460.5-461.8 m)	1,524.0- (464.5-	-
Oil shale, dusky- to dark-yellowish-brown, very thinly		
laminated. Pods and irregular lenses of moderate		8
brown translucent crystalline nahcolite	1,534.0-	
Nahcolite, pale-yellowish-brown, opaque, crystalline		-1,554.0 -473.7)
Oil shale, dusky-yellowish-brown, dusky brown,		
brownish-black, and minor dark-yellowish-brown,		
very thinly laminated, contains a .8-ft-thick		
(24 cm) nahcolite bed at 1,575.0 ft (480.I m)	1,554.0- (473.7-	
Nahcolite, white- to pale-yellowish-brown bedded,		
opaque, crystalline interbedded with laminae of		
oil shale	1,587.0- (483.7-	
Oil shale, like unit at 1,554-1,587 ft (473.7-483.7 m)	1,598.3- (487.2-	
No recovery		-1,600.8 -487.9)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, like unit at 1,554-1,587 ft (473.7-483.7 m)	1,600.8-1,603.8 (487.9-488.8)
Halite, bedded	1,603.8-1,604.3 (488.8-489.0)
Interbedded nahcolite and oil shale	1,604.3-1,610.0 (489.0-490.1)
Oil shale, like unit at 1,554-1,587 ft (473.7-483.7 m)	1,610.0-1,616.0 (490.1-492.6)
Interbedded nahcolite and oil shale	1,616.0-1,620.5 (492.6-493.9)
Oil shale, like unit at 1,554-1,587 ft (473.7-483.7 m)	1,620.5-1,622.6 (493.9-494.6)
Interbedded nahcolite and oil shale	1,622.6-1,629.0 (494.6-496.5)
Halite, bedded	1,629.0-1,629.5 (496.5-496.7)
Interbedded nahcolite and oil shale	1,629.5-1,632.0 (496.7-497.4)
No recovery (loss due to solubility of salts)	1,632.0-1,639.0 (497.4-499.6)
Oil shale, dusky-yellowish-brown, dark-yellowish-brown,	
and minor amounts of light brown, very thinly	
laminated. Interbedded with halite zones up to	
.5 ft (15 cm) thick, and moderate brown, crystalline	,
translucent nahcolite beds and "rosettes" some	
nahcolite is pale-yellowish-brown	1,639.0-1,660.0 (499.6-506.0)

Description	Interval thickness ft (m)	
CoreContinued .		
Oil shale, as above	1,660.0-1,667.7 (506.0-508.3)	
No recovery	1,667.7-1,669.0 (508.3-508.7)	
Oil shale, like unit at 1,639.0-1,660.0 ft		
(499.6-506.0 m) scattered nahcolite "rosettes."		
Base of R-5 zone at 1,680.9 ft (512.3 m)	1,669.0-1,680.9 (508.7-512.3)	
Nahcolite, dusky- to dark-yellowish-brown, contains a		
halite stringer at 1,682 ft (512.7 m)	1,680.9-1,686.9 (512.3-514.2)	
No recovery	1,686.9-1,699.0 (514.2-517.9)	
Nahcolite, like unit at 1,680.9-1,686.9 ft		
(512.3-514.2 m)	1,699.0-1,702.5 (517.9-518.9)	
Oil shale, dusky-yellowish-brown with minor amounts of		
pale yellowish brown, very thinly laminated.		
Moderate brown translucent, crystalline nahcolite		
"rosettes" up to 3 inches (8 cm) in diameter		
scattered throughout core	1,702.5-1,715.0 (518.9-522.7)	
Nahcolite, pale-yellowish-brown, opaque, crystalline,		
thin bedded	1,715.0-1,717.0 (522.7-523.3)	
Oil shale, like unit at 1,702.5-1,715.0 ft		
(518.9-522.7 m), contains some brownish-black		
oil shale	1,717.0-1,727.9 (523.3-526.7)	

Description	Interval thickness ft (m)
CoreContinued ,	
Oil shale, dark-yellowish-brown and dusky-yellowish-	
brown, very thinly laminated	1,768.5-1,777.5 (539.0-541.8)
Nahcolite, like unit at 1,759.0-1,768.5 ft	
(536.1-539.0 m)	1,777.5-1,779.5 (541.8-542.4)
Oil shale, like unit at 1,768.5-1,777.5 ft	3,
(539.0-541.8 m)	1,779.5-1,783.0 (542.4-543.5)
Nahcolite, like unit at 1,759.0-1,768.5 ft	
(536.Ī-539.0 m)	1,783.0-1,788.2 (543.5-545.0)
No recovery	1,788.2-1,789.0 (545.0-545.3)
Oil shale, like unit at 1,768.5-1,777.5 ft	
(539.0-541.8 m) with beds and "rosettes" (up to	
4 inches diameter, 10 cm) of nahcolite scattered	
throughout interval	1,789.0-1,818.9 (545.3-554.4)
No recovery	1,818.9-1,819.4 (554.4-554.6)
Oil shale, dusky-yellowish-brown, contains "rosettes"	
of nahcolite up to 9 inches (23 cm) in diameter	1,819.4-1,828.4 (554.6-557.3)
Nahcolite, bedded and "rosettes"	1,828.4-1,830.0 (557.3-557.3)

Description	Interval thickness ft (m)
CoreContinued	
No recovery	1,727.9-1,729.0 (526.7-527.0)
Oil shale, like unit at 1,717.0-1,727.9 ft	
(523.3-526.7)	1,729.0-1,734.5 (527.0-528.7)
Nahcolite, moderate-brown, translucent, crystalline,	6
thin bedded, with minor amounts of oil shale	1,734.5-1,737.0 (528.7-529.4)
Oil shale, like unit at 1,717.0-1,727.9 ft	
(523.3-526.7 m)	1,737.0-1,739.0 (529.4-530.0)
Nahcolite, like unit at 1,734.5-1,737.0 ft	
(528.7-529.4 m). Base of L-4 zone at 1,742 ft	
(531.0 m)	1,739.0-1,742.0 (530.0-531.0)
Oil shale, like unit at 1,717.0-1,727.9 ft	,
(523.3-526.7 m), contains a 4-inch-thick (10 cm)	
bed of nahcolite and halite at 1,750.9 ft	*
(533.7 m)	1,742.0-1,759.0 (531.0-536.1)
Nahcolite, pale-yellowish-brown and moderate brown,	
opaque to translucent, thin bedded with minor	
oil shale	1,759.0-1,768.5 (536.1-539.0)

Description	Interval thickness ft (m)
CoreContinued ,	
Interbedded and laminated oil shale, nahcolite and minor	n.
marlstone, pale yellowish brown to dusky yellowish	
brown. "Rosettes" up to 10 inches (25 cm) in	
diameter. Halite fills vertical fractures at	
1,834 and 1,846 ft (559.0 and 562.7 m)	1,830.0-1,850.0 (557.3-557.8)
Oil shale, dusky-yellowish-brown, laminated with dusky-	
to dark-yellowish-brown matlstone. Nahcolite	
"rosettes" up to 6 inches in diameter every foot	
or two. Fine- to medium-grained nahcolite bed at	
1,873-1,873.5 ft (570.9-571.0 m) and a .1-foot-	
thick (3 cm) bed at 1,853.8 ft (565.0 m)	1,850.0-1,874.0 (557.8-563.9)
No recovery	1,874.0-1,879.0 (563.9-572.7)
Nahcolite, interbedded with dark-yellowish-brown and	
minor dusky-yellowish-brown marlstone(?). Nahcolite	
beds range from .3 to 1.0 ft (8 to 30 cm) thick.	
Dark-gray siltstone layer at 1,887.5-1,887.7 ft	
(575.3-575.4 m). Nahcolite "rosettes" scattered	
throughout lower half of unit (1,893-1,907.4 ft,	
577.0-581.4 m)	1,879.0-1,907.4 (572.7-581.4)

Description	Interval thickness ft (m)
CoreContinued ,	
Oil shale, dark-yellowish-brown with minor laminae of	
dusky yellowish brown, thinly laminated. Bedded	1
opaque nahcolite at 1,908.2-1,908.4 and 1,921.3-	
1,921.5 ft (581.6-581.7 and 585.6-585.7 m).	
Nahcolite "rosettes" up to 7 inches (18 cm) in	
diameter, most are 3-5 inches (8-15 cm). Eroded	
"rosettes" cause core to be in .5-2 ft (15-60 cm)	
lengths	1,907.4-1,937.2 (581.4-590.5)
No recovery	1,937.2-1,937.4 (590.5-590.5+)
Oil shale, dark- to dusky-yellowish-brown, thinly	
laminated. Thin interbeds of nahcolite at 1,943,	
1,960, 1,961.5, 1,969.8-1,970.2, and 1,972.3-	
1,972.9 ft (592.2, 597.4, 597.8, 600.4-600.5, and	•
601.2-601.3 m). Nahcolite "rosettes" between	
1,946-1,967 ft (593.1-599.5 m) and scattered	š.
between 1,968-1,997 ft (599.8-608.7 m). Abundant	
dolomite lenses up to 1 inch (3 cm) long and	
1/8 inch (.3 cm) high throughout oil shale.	
"60 bed" 1,988-1,992 ft (605.9-607.2 m). Base	
of R-4 zone at 1,997 ft (608.7 m)	1,937.4-1,997.0 (590.5+-608.7)

Description	Interval thickness ft (m)
CoreContinued .	
Marlstone(?), dark-yellowish-brown, laminated with minor	
amounts of dusky yellowish brown. Unit contains	_
thin bedded nahcolite at 2,002.0, 2,004.6, 2,017.6,	
and 2,020.6 ft (610.2, 611.0, 615.0, and 615.9 m).	
Nahcolite "rosettes" 1-3 inches (3-8 cm) diameter	
below 2,015 ft (614.2 m). Bottom of L-3 zone at	-
2,026 ft (617.5 m)	1,997.0-2,025.9 (608.7-617.5)
No recovery	2,025.9-2,027.1 (617.5-617.9)
Oil shale, dark-yellowish-brown, interlaminated and	
interbedded with bedded nahcolite ranging from	
laminae to 3 inches (8 cm) thick. Unit contains	
pale-yellowish-brown lenses of dolomite up to	
2 inches (5 cm) long (most are 1/4-1 inch; 1-3 cm).	
Rock half or more nahcolite. "Rosettes" of	
nahcolite up to 1 1/2 ft (45 cm) in diameter	2,027.1-2,085.7 (617.9-635.7)
No recovery	2,085.7-2,087.0 (635.7-636.1)
Oil shale, like unit at 2,027.1-2,085.7 ft	
(617.9-635.7 m), with nahcolite beds up to	
6 inches (15 cm) thick	2,087.0-2,094.0 (636.1-638.3)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, dark- to dusky-yellowish-brown, containing	
nahcolite "rosettes"	2,094.0-2,116.3 (638.3-645.0)
No recovery	2,116.3-2,117.0 (645.0-645.3)
Nahcolite, dark-yellowish-brown, bedded	2,117.0-2,123.0 (645.3-647.1)
Oil shale, dusky- to dark-yellowish-brown; unit contains	
opaque and translucent brown nahcolite "rosettes"	
up to 1 ft (30 cm) in diameter (most are 3-5 inches,	
8-15 cm). Pale-yellow-brown lenses of dolomite	
below 2,137 ft (651.4 m)	2,123.0-2,146.5 (647.1-654.3)
No recovery	2,146.5-2,147.5 (654.3-654.6)
Oil shale, dusky-yellowish-brown interlaminated with	
minor dark-yellowish-brown marlstone. Upper 3 ft	
(90 cm) is interbedded marlstone, nahcolite, and	
nahcolite "rosettes." Base of R-3 zone	2,147.5-2,163.0 (654.6-659.3)
Marlstone(?) dark-yellowish-brown, thinly laminated	2,163.0-2,175.5 (659.3-663.1)
No recovery	2,175.5-2,177.5 (663.1-663.7)
Marlstone(?), like unit at 2,163-2,175.5 ft	
(659.3-663.1 m)	2,177.5-2,180.5 (663.7-664.6)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, dusky- and dark-yellowish-brown, thinly	
laminated	2,180.5-2,189.0 (664.6-667.2)
Nahcolite, bedded	2,189.0-2,195.0 (667.2-669.0)
Oil shale, dark- to dusky-yellowish-brown laminated;	*
minor amounts of marlstone(?) and bedded nahcolite.	
Nahcolite rosettes as large as 8 inches (20 cm)	
(most are 2-4 inches, 5-8 cm) scattered through-	
out core. Dolomite lenses in upper 32 ft (9.8 cm)	2,195.0-2,245.0 (669.0-684.3)
Marlstone(?), dark-yellowish-brown, laminated, minor	
amounts of nahcolite	2,245.0-2,254.3 (684.3-687.1)
No recovery	2,254.3-2,258.0 (687.1-688.2)
Oil shale, dusky- to dark-yellowish-brown, laminated	2,258.0-2,268.0 (688.2-691.3)
Oil shale(?), dark-yellowish-brown, laminated; contains	•
dolomite lenses 1/8-1/2 inch (.3-1 cm) long	2,268.0-2,281.0 (691.3-695.2)
No recovery. Blue marker at 2,284 ft (696.2 m); also	•
base of R-2 zone	2,281.0-2,286.0 (695.2-696.8)
Oil shale, dusky- to dark-yellowish-brown, laminated;	
dolomite lenses	2,286.0-2,290.5 (696.8-698.1)

Description	Interval thickness ft (m)
CoreContinued	
Oil shale, dusky-yellow-brown, laminated. Unit has	
what appear to be slickensides on horizontal	
fracture surfaces, probably the typical-appearing	
Garden Gulch member	2,290.5-2,299.0 (698.1-700.7)
Interbedded shale, marlstone and a white bedded	
material. Shale is bluish-gray from 2,305 to	
2,310 ft (702.6 to 704.1 m), marlstone is	
dusky-yellowish-brown, laminated to thin bedded;	
white material between 2,299 and 2,305 ft	
(700.7 and 702.6 m) in layers less than .1 ft	
(3 cm) thick. Core discs easily. Base of L-1	
zone	2,299.0-2,310.0 (700.7-704.1)
Interlaminated and interbedded, marlstone, oil shale	1 "
and shale, dark- to dusky-yellowish-brown.	
Fossils in 2,326-2,327-ft (709.0-709.3 m) interval,	
may be fish vertabrae. Whitish, very thin beds	
between 2,344.5-2,346.0 ft (714.6-715.1 m). Core	
discs easily. Core shattered in many places	2,310.0-2,382.0 (704.1-726.0)
No recovery	2,382.0-2,382.5 TT (726.0-726.2)